

UNIVERSITY OF ESWATINI



MAIN EXAMINATION 2019/2020

TITLE OF PAPER: ANALYTICAL CHEMISTRY II
COURSE NUMBER: CHE 411
TIME ALLOWED: THREE (3) HOURS
INSTRUCTIONS: ANSWER ANY FOUR (4) QUESTIONS

Special Requirements

1. Data sheet.

YOU ARE NOT SUPPOSED TO OPEN THIS PAPER UNTIL PERMISSION TO DO SO HAS BEEN GIVEN BY THE CHIEF INVIGILATOR.

QUESTION 1 [25]

- a) For the $\text{VO}^{2+}/\text{V}^{3+}$ system in acid, calculate the concentration of V^{3+} at $\text{pH}=3$ if the potential measured for a 0.0625M VO^{2+} solution is 0.562V vs SCE [5]
- b) Explain, using diagrams and equations, how the selectivity coefficient and ion exchange principles enable fabrication of a pNa electrode. [5]
- c) i) With the aid of a diagram, explain how a Saturated Calomel Electrode (SCE) is fabricated, and explain the role of each component in the electrode. [6]
- ii) Write down its half cell reaction and Nernst expression. [4]
- iii) Calculate the standard electrode potential for the SCE. [3]
- iv) Under what conditions will the SCE not work. [2]

QUESTION 2 [25]


- (a) Describe, giving examples of as well as drawings and half cell reactions of, indicator electrodes of [3]
- (i) The First Kind [3]
- (ii) The Second Kind [3]
- (iii) The Third Kind [3]
- (b) Write down the Nernst equation, and explain all the terms appearing in it. [3]
- (c) In potentiometry, potentials are measured relative to the standard hydrogen electrode (SHE) potential. [4]
- i) Draw the SHE, and label all its components [4]
- ii) Write down the electrochemical equation taking place within the SHE and state its standard electrode potential [2]
- (d) The data below were obtained when a F^- ion-selective electrode was immersed in a series of standard solutions whose ionic strength was constant at 2.0M .

$[\text{F}^-] (\text{M})$	$E (\text{mV})$
2.35×10^{-5}	-74.8
2.62×10^{-4}	-48.4
2.13×10^{-3}	-18.7
1.99×10^{-2}	-10.0
2.48×10^{-1}	+37.7

What is the concentration of F^- in the sample if it gave a reading of -22.5mV [7]

QUESTION 3 [25]

(a) For the electrochemical cell:



- i) What component is represented by the symbol “//”? [1]
- ii) Use a diagram to explain its role in potentiometry [3]
- iii) Use drawings to show how it is constructed [4]
- iv) Use illustrations to explain how it works [2]
- v) Calculate ΔG for this cell [5]
- vi) Would the cell described above be galvanic as written? [1]
- (b) Use equations to describe the role of Ti^{4+} intermediate in the coulometric titration of Fe^{3+} . [5]
- (c) Use diagrams and equations to describe how an amperometric titration of metal ion M^{2+} can be carried out with a one-polarized electrode system. [4]

QUESTION 4 [25]

- a) Indicator and reference electrodes are now combined in compact units to produce an instrument that measures voltages in electroanalytical instruments.
- i) With the aid of a diagram, use the ion exchange theory to explain how a compact pH glass membrane electrode works. [5]
- ii) Write the Nernst expression for an ideal pH glass electrode, and show that unit calibrations in the readout are in increments of 59mV per decade change of H^+ concentration. [5]
- iii) Explain, using diagrams and equations, how the selectivity coefficient and ion exchange principles enable fabrication of a pNa electrode. [5]
- b) i) Outline the steps involved in the calibration of pH glass electrodes. [3]
- ii) List two (2) sources of standards used in the calibration of pH glass electrodes. [2]
- c) It takes 9.085 minutes to titrate a 10.053g sample of Na_2CO_3 coulometrically in an electrolytic cell with electrogenerated hydrogen ions. The generating current is 205.16 mA in a system incorporating Pt electrodes. Assuming that the endpoint occurs when all CO_3^{2-} has been converted to H_2CO_3 , calculate the concentration Na_2CO_3 in the sample in % units. [5]

QUESTION 5 [25]

- a) Describe the term "overpotential" in relation to the polarography technique, and explain why overpotential is desirable in this electroanalytical technique. [3]
- b) Draw and label the electrode used in classical polarography and explain how it works. [6]
- c) i) Explain why inert electrolyte such as KNO_3 is added to solutions in large quantities prior to measurement by classical polarography [2]
 ii) Explain why nitrogen gas is bubbled through aqueous solutions prior to polarographic measurements [2]
- d) For each of the following techniques, indicate, on a voltage-time plot, when sampling of the signal is carried out. Draw the shape of the resultant voltammogram, and indicate the typical resolution (in Volts) and detection limit (in mol/L). [4]
 i) Tast Polarography. [4]
 ii) Normal Pulse Polarography. [4]
 iii) Differential Pulse Polarography [4]

QUESTION 6 [25]

- a) i) Use a diagram to illustrate the dependence of "non-faradaic" current on time during the lifetime of a mercury drop in polarography. [3]
 ii) Use a diagram to illustrate the dependence of "faradaic" current on time during the lifetime of a mercury drop in polarography. [3]
 iii) Use a diagram to illustrate the effect of concentration on "non-faradaic" current during the lifetime of a mercury drop in polarography. [3]
- b) A solution of 0.200M Cu^{2+} in 1M H^+ , resistance $0.5\ \Omega$, is to be electrodeposited to 99.995% completion with 1A in an open cell (partial pressure of O_2 in air = 0.2 atm). In the equation $E_{\text{app}} = E_{\text{cathode}} + IR + \mathcal{Q}$ used to ascertain the potential at which electrodeposition will occur:
 i) Calculate E_{cathode} . [1]
 ii) Calculate E_{anode} . [1]
 iii) Calculate the IR drop. [1]
 iv) Describe the term \mathcal{Q} , and explain its origins in electrogravimetry using suitable equations. [3]
- a) Consider the voltametric titration of Tl^+ with electrochemically generated Br_2 according to the reaction
- $$\text{Tl}^+ + \text{Br}_2 \rightleftharpoons \text{Tl}^{3+} + 2\text{Br}^-, \text{ where}$$
- $$\text{Tl}^{3+} + 2\text{e}^- \rightleftharpoons \text{Tl}^+ \quad E^0 = -0.78\text{V vs SCE}$$
- $$\text{Br}_2 + 2\text{e}^- \rightleftharpoons 2\text{Br}^- \quad E^0 = -1.08\text{V vs SCE}$$
- i) Draw the current-voltage curves of this titration at the following stages of the titration: [8]
 $f = 0;$ $f = 0.5;$ $f = 1.0;$ $f = 1.5$
- ii) Plot the titration curve expected for this system using a single indicator electrode. [2]

PERIODIC TABLE OF ELEMENTS

GROUPS

PERIODS	GROUPS																	
	IA	IIA	III	IV	V	VI	VII	VIII	IX	X	XI	XII	XIII	XIV	XV	XVI	XVII	XVIII
1	H 1.008																	He 4.003
2	Li 6.941	Be 9.012											B 10.811	C 12.011	N 14.007	O 15.999	F 18.998	Ne 20.180
3	Na 22.990	Mg 24.305											Al 26.982	Si 28.086	P 30.974	S 32.06	Cl 35.453	Ar 39.948
4	K 39.098	Ca 40.078	Sc 44.956	Ti 47.88	V 50.942	Cr 51.996	Mn 54.938	Fe 55.847	Ni 58.69	Cu 63.546	Zn 65.39	Ga 69.723	Ge 72.61	As 74.922	Se 78.96	Br 79.904	Kr 83.80	
5	Rb 85.468	Sr 87.62	Y 88.906	Zr 91.224	Nb 92.906	Mo 95.94	Tc 98.907	Ru 101.07	Rh 106.42	Pd 107.87	Cd 112.41	In 114.82	Sn 118.71	Sb 121.75	Te 127.60	I 126.90	Xe 131.29	
6	Cs 132.91	Ba 137.33	*La 138.91	Hf 178.49	Ta 180.95	W 183.85	Re 186.21	Os 190.2	Pt 195.08	Au 196.97	Hg 200.59	Tl 204.38	Pb 207.2	Bi 208.98	Po 209	At 210	Rn 222	
7	Fr 223	Ra 226.03	**Ac 227	Rf 261	Ha 262	Unh 263	Uns 267	Uno 269	Uue 269	Uuh 271	Uuq 272	Uur 273	Uus 277	Uut 281	Uuq 285	Uuo 286	Uuq 287	

Atomic mass
Symbol
Atomic No.

TRANSITION ELEMENTS

140.12	Ce 58	Pr 59	Nd 60	Pm 61	Sm 62	Eu 63	Gd 64	Tb 65	Dy 66	Ho 67	Er 68	167.26	168.93	174.97
232.04	Th 90	Pa 91	U 92	Np 93	Pu 94	Am 95	Cm 96	Bk 97	Cf 98	Es 99	Fm 100	(257)	(258)	(260)

* Lanthanide Series

** Actinide Series

() indicates the mass number of the isotope with the longest half-life.

Quantity	Symbol	Value	General data and fundamental constants
Speed of light	c	$2.997\,924\,58 \times 10^8 \text{ m s}^{-1}$	
Elementary charge	e	$1.602\,177 \times 10^{-19} \text{ C}$	
Faraday constant	$F = eN_A$	$9.6485 \times 10^4 \text{ C mol}^{-1}$	
Boltzmann constant	k	$1.380\,65 \times 10^{-23} \text{ J K}^{-1}$	
Gas constant	$R = kN_A$	$8.314\,5 \text{ J K}^{-1} \text{ mol}^{-1}$ $8.205\,78 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$	
Planck constant	h	$6.626\,08 \times 10^{-34} \text{ J s}$	
	$\hbar = h/2\pi$	$1.054\,57 \times 10^{-34} \text{ J s}$	
Avogadro constant	N_A	$6.022\,14 \times 10^{23} \text{ mol}^{-1}$	
Atomic mass unit	u	$1.660\,54 \times 10^{-27} \text{ kg}$	
Mass of electron	m_e	$9.109\,39 \times 10^{-31} \text{ kg}$	
Proton	m_p	$1.672\,62 \times 10^{-27} \text{ kg}$	
Neutron	m_n	$1.674\,93 \times 10^{-27} \text{ kg}$	
Vacuum permeability	μ_0	$4\pi \times 10^{-7} \text{ J A}^{-2} \text{ C}^{-2} \text{ m}^{-1}$	
Vacuum permittivity	$\epsilon_0 = 1/c^2 \mu_0$	$8.854\,18 \times 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$	
	$\epsilon = \epsilon_0$	$1.112\,65 \times 10^{-10} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$	
Sohr magneton	$\mu_B = e\hbar/2m_e$	$9.274\,02 \times 10^{-24} \text{ J T}^{-1}$	
Nuclear magneton	$\mu_N = e\hbar/2m_p$	$5.050\,78 \times 10^{-27} \text{ J T}^{-1}$	
Electron g value	g_e	2.002 31	
Sohr radius	$a_0 = 4\pi\epsilon_0\hbar^2/m_e e^2$	$5.291\,77 \times 10^{-11} \text{ m}$	
Ryberg constant	$R_\infty = m_e e^4/8h^3\epsilon_0$	$1.097\,37 \times 10^8 \text{ m}^{-1}$	
Fine structure constant	$\alpha = \mu_0 e^2 c/2h$	$7.297\,35 \times 10^{-2}$	
Gravitational constant	G	$6.672\,59 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$	
Standard acceleration of free fall	g	$9.806\,65 \text{ m s}^{-2}$	

† Exact (defined) values

f	p	n	μ	m	c	d	k	M	G	Prefixes
femto	pico	nano	micro	milli	centi	deci	kilo	mega	giga	
10^{-15}	10^{-12}	10^{-9}	10^{-6}	10^{-3}	10^{-2}	10^{-1}	10^1	10^6	10^9	

APPENDIX C (continued)

Half-reaction	E° (V)
$\text{Hg}_2\text{Cl}_2(s) + 2e^-$	$= 2\text{Hg} + 2\text{Cl}^-$ 0.2676
$\text{BiO}^+ + 2\text{H}^+ + 3e^-$	$= \text{Bi} + \text{H}_2\text{O}$ 0.32
$\text{AgCl}(s) + e^-$	$= \text{Ag} + \text{Cl}^-$ 0.2221
$\text{SbO}^+ + 2\text{H}^+ + 3e^-$	$= \text{Sb} + \text{H}_2\text{O}$ 0.212
$\text{CuCl}_2 + e^-$	$= \text{Cu} + 2\text{Cl}^-$ 0.178
$\text{SO}_4^{2-} + 4\text{H}^+ + 2e^-$	$= \text{SO}_2(g) + 2\text{H}_2\text{O}$ 0.17
$\text{Sn}^{4+} + 2e^-$	$= \text{Sn}^{2+}$ 0.15
$\text{S} + 2\text{H}^+ + 2e^-$	$= \text{H}_2\text{S}(g)$ 0.14
$\text{TiO}^{2+} + 2\text{H}^+ + e^-$	$= \text{Ti}^{3+} + \text{H}_2\text{O}$ 0.10
$\text{S}_4\text{O}_6^{2-} + 2e^-$	$= 2\text{S}_2\text{O}_3^{2-}$ 0.08
$\text{AgBr}(s) + e^-$	$= \text{Ag} + \text{Br}^-$ 0.071
$2\text{H}^+ + 2e^-$	$= \text{H}_2$ 0.0000
$\text{Pb}^{2+} + 2e^-$	$= \text{Pb}$ -0.126
$\text{Sn}^{2+} + 2e^-$	$= \text{Sn}$ -0.136
$\text{AgI}(s) + e^-$	$= \text{Ag} + \text{I}^-$ -0.152
$\text{Mo}^{3+} + 3e^-$	$= \text{Mo}$ approx -0.2
$\text{N}_2 + 5\text{H}^+ + 4e^-$	$= \text{H}_2\text{NNH}_3^+$ -0.23
$\text{Ni}^{2+} + 2e^-$	$= \text{Ni}$ -0.246
$\text{V}^{2+} + e^-$	$= \text{V}^{3+}$ -0.255
$\text{Co}^{2+} + 2e^-$	$= \text{Co}$ -0.277
$\text{Ag}(\text{CN})_2^- + e^-$	$= \text{Ag} + 2\text{CN}^-$ -0.31
$\text{Cd}^{2+} + 2e^-$	$= \text{Cd}$ -0.403
$\text{Cr}^{3+} + e^-$	$= \text{Cr}^{2+}$ -0.41
$\text{Fe}^{3+} + e^-$	$= \text{Fe}^{2+}$ -0.440
$2\text{CO}_2 + 2\text{H}^+ + 2e^-$	$= \text{H}_2\text{C}_2\text{O}_4$ -0.49
$\text{H}_3\text{PO}_3 + 2\text{H}^+ + 2e^-$	$= \text{H}_2\text{P}_2\text{O}_5 + \text{H}_2\text{O}$ -0.50
$\text{U}^{4+} + e^-$	$= \text{U}^{3+}$ -0.61
$\text{Zn}^{2+} + 2e^-$	$= \text{Zn}$ -0.763
$\text{Cr}^{3+} + 3e^-$	$= \text{Cr}$ -0.71
$\text{Mn}^{2+} + 2e^-$	$= \text{Mn}$ -1.18
$\text{Zr}^{4+} + 4e^-$	$= \text{Zr}$ -1.53
$\text{Ti}^{3+} + 3e^-$	$= \text{Ti}$ -1.63
$\text{Al}^{3+} + 3e^-$	$= \text{Al}$ -1.66
$\text{Th}^{4+} + 4e^-$	$= \text{Th}$ -1.90
$\text{Mg}^{2+} + 2e^-$	$= \text{Mg}$ -2.37
$\text{La}^{3+} + 3e^-$	$= \text{La}$ -2.52
$\text{Na}^+ + e^-$	$= \text{Na}$ -2.714
$\text{Ca}^{2+} + 2e^-$	$= \text{Ca}$ -2.87
$\text{Sr}^{2+} + 2e^-$	$= \text{Sr}$ -2.89
$\text{K}^+ + e^-$	$= \text{K}$ -2.925
$\text{Li}^+ + e^-$	$= \text{Li}$ -3.045

APPENDIX C. POTENTIALS OF SELECTED HALF REACTIONS AT 25 °C

A summary of oxidation/reduction half-reactions arranged in order of decreasing oxidation strength and useful for selecting reagent systems.

Half-reaction	E° (V)
$F_2(g) + 2H^+ + 2e^- = 2HF$	3.06
$O_2 + 2H^+ + 2e^- = O_2 + H_2O$	2.07
$S_2O_8^{2-} + 2e^- = 2SO_4^{2-}$	2.01
$Ag^+ + e^- = Ag$	2.00
$H_2O_2 + 2H^+ + 2e^- = 2H_2O$	1.77
$MnO_4^- + 4H^+ + 3e^- = MnO_2(s) + 2H_2O$	1.70
$Ce(IV) + e^- = Ce(III) \text{ (in 1M HClO}_4\text{)}$	1.61
$H_2IO_6 + H^+ + 2e^- = IO_3^- + 3H_2O$	1.6
$BiO_3^- \text{ (biomartite)} + 4H^+ + 2e^- = 2BiO^+ + 2H_2O$	1.59
$BrO_3^- + 6H^+ + 5e^- = \frac{1}{2}Br_2 + 3H_2O$	1.52
$MnO_4^- + 6H^+ + 5e^- = Mn^{2+} + 4H_2O$	1.51
$PbO_2 + 4H^+ + 2e^- = Pb^{2+} + 2H_2O$	1.455
$Cl_2 + 2e^- = 2Cl^-$	1.36
$Cr_2O_7^{2-} + 14H^+ + 6e^- = 2Cr^{3+} + 7H_2O$	1.33
$MnO_4^-(aq) + 4H^+ + 2e^- = Mn^{2+} + 2H_2O$	1.23
$O_2(g) + 4H^+ + 4e^- = 2H_2O$	1.229
$IO_3^- + 6H^+ + 5e^- = \frac{1}{2}I_2 + 3H_2O$	1.20
$Br_2(l) + 2e^- = 2Br^-$	1.055
$IO_3^- + e^- = \frac{1}{2}I_2 + 2Cl^-$	1.05
$VO_2^+ + 2H^+ + e^- = VO^{2+} + H_2O$	1.00
$HNO_3 + H^+ + e^- = NO(g) + H_2O$	1.00
$NO_3^- + 3H^+ + 2e^- = HNO_2 + H_2O$	0.94
$2Hg^{2+} + 2e^- = Hg_2^{2+}$	0.92
$Cu^+ + e^- = Cu(s)$	0.86
$Ag^+ + e^- = Ag$	0.799
$Hg_2^{2+} + 2e^- = 2Hg$	0.79
$Fe^{3+} + e^- = Fe^{2+}$	0.771
$O_2(g) + 2H^+ + 2e^- = H_2O_2$	0.682
$2HgCl_2 + 2e^- = Hg_2Cl_2(l) + 2Cl^-$	0.63
$Hg_2SO_4(s) + 2e^- = 2Hg + SO_4^{2-}$	0.615
$Sb_2O_5 + 6H^+ + 4e^- = 2Sb^{3+} + 3H_2O$	0.581
$H_2AsO_4 + 2H^+ + 2e^- = HAsO_2 + 2H_2O$	0.559
$I_2 + 2e^- = 2I^-$	0.545
$Cu^+ + e^- = Cu$	0.52
$VO^{2+} + 2H^+ + e^- = V^{3+} + H_2O$	0.337
$Fe(CN)_6^{3-} + e^- = Fe(CN)_6^{4-}$	0.35
$Cu^{2+} + e^- = Cu$	0.317
$UO_2^{2+} + 4H^+ + 2e^- = U^{4+} + 2H_2O$	0.334

(continued)